



MEDICATION GUIDE
Sumatriptan and Naproxen Sodium Tablets
(Soo-ma-TRIP-tan and na-PROX-en SOE-dee-um)

Read this Medication Guide before you start taking sumatriptan and naproxen sodium tablets and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking with your healthcare provider about your medical condition or treatment.

What is the most important information I should know about sumatriptan and naproxen sodium tablets?
Sumatriptan and naproxen sodium tablets may increase your chance of a heart attack or stroke that can lead to death. Sumatriptan and naproxen sodium contain 2 medicines: sumatriptan and naproxen sodium (a nonsteroidal anti-inflammatory drug [NSAID]).

- This risk may happen early in treatment and may increase:
 - with increasing doses of NSAIDs
 - with longer use of NSAIDs

Do not take sumatriptan and naproxen sodium tablets right before or after a heart surgery called a "coronary artery bypass graft (CABG)."

Avoid taking sumatriptan and naproxen sodium tablets after a recent heart attack, unless your healthcare provider tells you to. You may have an increased risk of another heart attack if you take NSAIDs after a recent heart attack.

Stop taking sumatriptan and naproxen sodium tablets and get emergency help right away if you have any of the following symptoms of a heart attack or stroke:

- discomfort in the center of your chest that lasts for more than a few minutes, or that goes away and comes back
- severe tightness, pain, pressure, or heaviness in your chest, throat, neck, or jaw
- pain or discomfort in your arms, back, neck, jaw, or stomach
- shortness of breath with or without chest discomfort
- breaking out in a cold sweat
- nausea or vomiting
- feeling lightheaded
- weakness in one part or on one side of your body
- slurred speech

Sumatriptan and naproxen sodium tablets are not for people with risk factors for heart disease unless a heart exam is done and shows no problem. You have a higher risk for heart disease if you:

- have high blood pressure
- have high cholesterol levels
- smoke
- are overweight
- have diabetes
- have a family history of heart disease

Sumatriptan and naproxen sodium tablets can cause ulcers and bleeding in the stomach and intestines at any time during your treatment.

Ulcers and bleeding can happen without warning symptoms and may cause death.

Your chance of getting an ulcer or bleeding increases with:

- past history of stomach ulcers, or stomach or intestinal bleeding with use of NSAIDs
- the use of medicines called "corticosteroids," "anticoagulants," and antidepressant medicines called "SSRIs" or "SNRIs"
- longer use
- more frequent use
- smoking
- drinking alcohol
- older age
- having poor health
- advanced liver disease
- bleeding problems

Sumatriptan and naproxen sodium tablets may cause serious allergic reactions or serious skin reactions that can be life-threatening. Stop taking sumatriptan and naproxen sodium tablets and get emergency help right away if you develop:

- sudden wheezing
- swelling of your lips, tongue, throat or body
- rash
- fainting
- problems breathing or swallowing
- reddening of your skin with blisters or peeling
- blisters or bleeding of your lips, eye lids, mouth, nose, or genitals

Sumatriptan and naproxen sodium tablets should only be used exactly as prescribed, at the lowest dose possible for your treatment, and for the shortest time needed.

Sumatriptan and naproxen sodium tablets already contain an NSAID (naproxen). Do not use sumatriptan and naproxen sodium tablets with other medicines to lessen pain or fever or with other medicines for colds or sleeping problems without talking to your healthcare provider first, because they may contain an NSAID also.

What are Sumatriptan and Naproxen Sodium Tablets?

Sumatriptan and naproxen sodium tablets are a prescription medicine that contains sumatriptan and naproxen sodium (an NSAID). Sumatriptan and naproxen sodium tablets are used to treat acute migraine headaches with or without aura in patients 12 years of age and older.

Sumatriptan and naproxen sodium tablets are not used to treat other types of headaches such as hemiplegic (that make you unable to move on one side of your body) or basilar (rare form of migraine with aura) migraines.

Sumatriptan and naproxen sodium tablets are not used to prevent or decrease the number of migraine headaches you have.

It is not known if sumatriptan and naproxen sodium tablets are safe and effective to treat cluster headaches.

Who should not take sumatriptan and naproxen sodium tablets?

Do not take sumatriptan and naproxen sodium tablets if you have:

- heart problems, history of heart problems, or right before or after heart bypass surgery
- had a stroke, transient ischemic attack (TIAs), or problems with your blood circulation
- hemiplegic migraines or basilar migraines. If you are not sure if you have these types of migraines, ask your healthcare provider.
- narrowing of blood vessels to your legs and arms (peripheral vascular disease), stomach (ischemic bowel disease), or kidneys
- uncontrolled high blood pressure
- taken any medicines in the last 24 hours that are called 5-HT₁ agonists that are triptans or contain ergotamine. Ask your healthcare provider for a list of these medicines if you are not sure.
- taken an antidepressant medicine called a monoamine oxidase (MAO) inhibitor within the last 2 weeks. Ask your healthcare provider for a list if you are not sure.
- had an asthma attack, hives, or other allergic reaction with aspirin or any other NSAID medicine
- an allergy to sumatriptan, naproxen, or any of the ingredients in sumatriptan and naproxen sodium tablets. See "What are the ingredients in sumatriptan and naproxen sodium tablets?" below for a complete list of ingredients.
- liver problems

What should I tell my healthcare provider before taking sumatriptan and naproxen sodium tablets?

Before you take sumatriptan and naproxen sodium tablets, tell your healthcare provider about all of your medical conditions, including if you:

- have high blood pressure
- have asthma
- have high cholesterol
- have diabetes
- smoke
- are overweight
- have heart problems or a family history of heart problems or stroke
- have kidney problems
- have liver problems
- have had epilepsy or seizures
- are not using effective birth control
- are pregnant, think you might be pregnant, or are trying to become pregnant. Taking NSAIDs, including sumatriptan and naproxen sodium tablets, at about 20 weeks of pregnancy or later may harm your unborn baby. If you need to take NSAIDs for more than 2 days when you are between 20 and 30 weeks of pregnancy, your healthcare provider may need to monitor the amount of fluid in your womb around your baby. **You should not take NSAIDs after about 30 weeks of pregnancy.**
- are breastfeeding or plan to breastfeed. The components of sumatriptan and naproxen sodium tablets pass into your breast milk and may harm your baby. Talk with your healthcare provider about the best way to feed your baby if you take sumatriptan and naproxen sodium tablets.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Sumatriptan and naproxen sodium tablets and certain other medicines can affect each other, causing serious side effects.

How should I take sumatriptan and naproxen sodium tablets?

- Certain people should take their first dose of sumatriptan and naproxen sodium tablets in their healthcare provider's office or in another medical setting. Ask your healthcare provider if you should take your first dose in a medical setting.
- Take sumatriptan and naproxen sodium tablets exactly as your healthcare provider tells you to take them.
- Take sumatriptan and naproxen sodium tablets whole with water or other liquids.
- Sumatriptan and naproxen sodium tablets can be taken with or without food.
- If you do not get any relief after your first dose, do not take a second dose without first talking with your healthcare provider.

Folding:

30 pages Pad

365 ---2--91.25 mm

235---0--235 mm

Glueing on 235 mm side

365 mm

Size: 365x235 mm



Sumatriptan & Naproxen Sodium Tabs 85 mg/500mg

Artwork Type: **MEDICATION GUIDE**

Artwork Code: **5220701**

Void Code: **5204021 (PGPI0363A)**

Dimension: **365x235 mm**

Void A/W Reason: **Revision is in line to the RLD PI (SLC changes) approved on April 28, 2021**

Country: **ANDA-US (NORTHSTAR)**

Language: **ENGLISH**

Mfg. Location: **DADRA**

Specification/Type of Paper: **Super Fine 41 GSM ITC Paper**

Folding:

30 pages Pad

365 ---2--91.25 mm

235---0--235 mm

Glueing on 235 mm side

Special Req.: –

Remark (if any):

Prepared by: **SAPNA**

Approved by RA:

APPROVAL HISTORY ATTACHED

No. of Colors: 1

Black

- If your headache comes back or you only get some relief from your headache:
 - For adults: a second dose may be taken 2 hours after the first dose. Do not take more than 2 doses of sumatriptan and naproxen sodium tablets 85 mg/500 mg in a 24-hour period.
 - For children 12 to 17 years of age: it is not known if taking more than 1 dose of sumatriptan and naproxen sodium tablets in 24 hours is safe and effective. Talk to your healthcare provider about what to do if your headache does not go away or comes back.
- If you take too many sumatriptan and naproxen sodium tablets, call your healthcare provider or go to the nearest hospital emergency room right away.
- You should write down when you have headaches and when you take sumatriptan and naproxen sodium tablets so you can talk with your healthcare provider about how sumatriptan and naproxen sodium tablets are working for you.

What should I avoid while taking sumatriptan and naproxen sodium tablets?

Sumatriptan and naproxen sodium tablets can cause dizziness, weakness, or drowsiness. If you have these symptoms, do not drive a car, use machinery, or do anything where you need to be alert.

What are the possible side effects of sumatriptan and naproxen sodium tablets?

Sumatriptan and naproxen sodium tablets may cause serious side effects. See "What is the most important information I should know about sumatriptan and naproxen sodium tablets?"

These serious side effects include:

- changes in color or sensation in your fingers and toes (Raynaud's syndrome)
- new or worse high blood pressure
- heart failure from body swelling (fluid retention)
- kidney problems including kidney failure
- low red blood cells (anemia)
- liver problems including liver failure
- asthma attacks in people who have asthma
- stomach and intestinal problems (gastrointestinal and colonic ischemic events). Symptoms of gastrointestinal and colonic ischemic events include:
 - sudden or severe stomach pain
 - stomach pain after meals
 - weight loss
 - nausea or vomiting
 - constipation or diarrhea
 - bloody diarrhea
 - fever

- problems with blood circulation to your legs and feet (peripheral vascular ischemia). Symptoms of peripheral vascular ischemia include:
 - cramping and pain in your legs or hips
 - feeling of heaviness or tightness in your leg muscles
 - burning or aching pain in your feet or toes while resting
 - numbness, tingling, or weakness in your legs
 - cold feeling or color changes in 1 or both legs or feet
- medication overuse headaches. Some people who use too many sumatriptan and naproxen sodium tablets may have worse headaches (medication overuse headache). If your headaches get worse, your healthcare provider may decide to stop your treatment with sumatriptan and naproxen sodium tablets.
- serotonin syndrome. Serotonin syndrome is a rare but serious problem that can happen in people using sumatriptan and naproxen sodium tablets, especially if sumatriptan and naproxen sodium tablets are used with antidepressant medicines called SSRIs or SNRIs. Stop taking sumatriptan and naproxen sodium tablets and call your healthcare provider right away if you have any of the following symptoms of serotonin syndrome:
 - changes in blood pressure
 - fast heartbeat
 - tight muscles
 - high body temperature
 - mental changes such as seeing things that are not there (hallucinations), agitation, or coma
 - trouble walking
 - seizures. Seizures have happened in people taking sumatriptan, one of the ingredients in sumatriptan and naproxen sodium tablets, who have never had seizures before. Talk with your healthcare provider about your chance of having seizures while you take sumatriptan and naproxen sodium tablets.

The most common side effects of sumatriptan and naproxen sodium tablets include:

- dizziness
- feeling weak, drowsy, or tired
- pain, discomfort, or stiffness in your neck, throat, jaw, or chest
- nausea
- tingling or numbness in your fingers or toes
- heartburn
- dry mouth
- feeling hot
- heartbeat problems
- muscle tightness

Stop sumatriptan and naproxen sodium tablets and call your healthcare provider right away if you have any of the following symptoms:

- nausea that seems out of proportion to your migraine
- sudden or severe stomach pain
- vomit blood
- blood in your bowel movement or it is black and sticky like tar
- yellow skin or eyes
- unusual weight gain
- more tired or weaker than usual
- flu-like symptoms
- itching
- diarrhea
- swelling of the arms, legs, hands, and feet
- tenderness in your upper right side

Tell your healthcare provider if you have any side effects that bother you or do not go away.

These are not all of the side effects of sumatriptan and naproxen sodium tablets. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store sumatriptan and naproxen sodium tablets?

Store sumatriptan and naproxen sodium tablets at room temperature between 68°F to 77°F (20°C to 25°C).

Keep sumatriptan and naproxen sodium tablets and all medicines out of the reach of children.

General information about the safe and effective use of sumatriptan and naproxen sodium tablets

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use sumatriptan and naproxen sodium tablets for a condition for which it was not prescribed. Do not give sumatriptan and naproxen sodium tablets to other people, even if they have the same problem you have. It may harm them.

This Medication Guide summarizes the most important information about sumatriptan and naproxen sodium tablets. If you would like more information, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about sumatriptan and naproxen sodium tablets that is written for healthcare professionals.

For more information, call Northstar Rx LLC at 1-800-206-7821

What are the ingredients in sumatriptan and naproxen sodium tablets?

Active ingredients: sumatriptan succinate and naproxen sodium.

Inactive ingredients: microcrystalline cellulose, dibasic calcium phosphate anhydrous, hypromellose, povidone, croscarmellose sodium, colloidal silicon dioxide, sodium chloride, sodium citrate, crospovidone, magnesium stearate, titanium dioxide, polyethylene glycol, FD&C Blue No. 2, FD&C Yellow No. 6, and iron oxide yellow.

Manufactured for:
Northstar Rx LLC
Memphis, TN 38141.

Manufactured by:
Sun Pharmaceutical Industries Ltd.
Survey No. 259/15, Dadra-396 191,
(U.T. of D & NH), India.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Medication Guides available at
www.northstarrxllc.com/products or call 1-800-206-7821.

5220701
Rev 06/2021

Folding:

30 pages Pad

365 ---2--91.25 mm

235---0--235 mm

Glueing on 235 mm side

365 mm

Size: 365x235 mm